



6% CopperHead® ROL

A Free Choice Mineral for Cattle and Horses

For control of fecal flies in manure of treated cattle and horses. Prevents development of face, horn, house and stable flies in the manure of treated cattle. Prevents development of house and stable flies in the manure of treated horses.

ACTIVE INGREDIENT:

| | |
|---|----------------|
| Tetrachlorvinphos* (CAS # 22248-79-9) | 0.494% |
| OTHER INGREDIENTS** | 99.506% |
| Total | 100.000% |

*RABON® Insecticide - Contains 2.24 grams RABON® per pound

**Refers only to ingredients which are not larvicidal

Guaranteed Analysis

| | |
|-------------------------|---------------|
| Calcium (Min) | 19.00% |
| Calcium (Max) | 22.00% |
| Phosphorus (Min) | 6.00% |
| Salt (Min) | 12.50% |
| Salt (Max) | 15.00% |
| Magnesium (Min) | 2.00% |
| Cobalt (Min) | 15 ppm |
| Copper (Min) | 1,800 ppm |
| Iodine (Min) | 60 ppm |
| Manganese (Min) | 5,400 ppm |
| Selenium (Min) | 16 ppm |
| Zinc (Min) | 5,400 ppm |
| Vitamin A (Min) | 165,000 IU/lb |
| Vitamin D-3 (Min) | 40,000 IU/lb |
| Vitamin E (Min) | 200 IU/lb |

Ingredients

Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Processed Grain By-Products, Magnesium Oxide, Molasses Products, Roughage Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Manganese Protienate, Zinc Oxide, Zinc Sulfate, Zinc Proteinate, Copper Sulfate, Copper Proteinate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Cobalt Proteinate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

Fly Control Program

SWEETLIX® 6% CopperHead® ROL prevents development of fecal flies in the manure of treated cattle and horses, but is not effective against existing adult flies. Use other control measures to reduce the existing adult fly population. In order to achieve optimum fly control, use **SWEETLIX® 6% CopperHead® ROL** in conjunction with other good management and sanitation practices. Start feeding **SWEETLIX® 6% CopperHead® ROL** in early spring before flies begin to appear and continue feeding throughout the year until cold weather restricts fly activity. **SWEETLIX® 6% CopperHead® ROL** may be fed to cattle up until slaughter and to lactating cows.

KEEP OUT OF REACH OF CHILDREN

CAUTION

(See Reverse for First Aid, Precautionary Statements and complete Directions for Use.)

NET WEIGHT: 50 LBS. (22.7 KG)

EPA REG NO. 7698-27

EPA EST. NO. 7698-AL-001

Rabon® is a registered trademark of Bayer HealthCare LLC



7 4 8 9 3 6 0 2 6 9 8 4

Batch #

10897

1118

Manufactured By: **Ridley USA Inc.**
111 West Cherry Street, Suite 500
Mankato, MN 56001



6% CopperHead® ROL

A Free Choice Mineral for Cattle and Horses

For control of fecal flies in manure of treated cattle and horses. Prevents development of face, horn, house and stable flies in the manure of treated cattle. Prevents development of house and stable flies in the manure of treated horses.

ACTIVE INGREDIENT:

| | |
|---|----------------|
| Tetrachlorvinphos* (CAS # 22248-79-9) | 0.494% |
| OTHER INGREDIENTS** | 99.506% |
| Total | 100.000% |

*RABON® Insecticide - Contains 2.24 grams RABON® per pound

**Refers only to ingredients which are not larvicidal

Guaranteed Analysis

| | |
|-------------------------|---------------|
| Calcium (Min) | 19.00% |
| Calcium (Max) | 22.00% |
| Phosphorus (Min) | 6.00% |
| Salt (Min) | 12.50% |
| Salt (Max) | 15.00% |
| Magnesium (Min) | 2.00% |
| Cobalt (Min) | 15 ppm |
| Copper (Min) | 1,800 ppm |
| Iodine (Min) | 60 ppm |
| Manganese (Min) | 5,400 ppm |
| Selenium (Min) | 16 ppm |
| Zinc (Min) | 5,400 ppm |
| Vitamin A (Min) | 165,000 IU/lb |
| Vitamin D-3 (Min) | 40,000 IU/lb |
| Vitamin E (Min) | 200 IU/lb |

Ingredients

Calcium Carbonate, Monocalcium Phosphate, Dicalcium Phosphate, Salt, Processed Grain By-Products, Magnesium Oxide, Molasses Products, Roughage Products, Hydrogenated Soybean Oil, Manganous Oxide, Manganese Sulfate, Manganese Protienate, Zinc Oxide, Zinc Sulfate, Zinc Proteinate, Copper Sulfate, Copper Proteinate, Sodium Selenite, Ethylenediamine Dihydroiodide, Calcium Iodate, Cobalt Carbonate, Cobalt Proteinate, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Mineral Oil and Red Iron Oxide.

Fly Control Program

SWEETLIX® 6% CopperHead® ROL prevents development of fecal flies in the manure of treated cattle and horses, but is not effective against existing adult flies. Use other control measures to reduce the existing adult fly population. In order to achieve optimum fly control, use **SWEETLIX® 6% CopperHead® ROL** in conjunction with other good management and sanitation practices. Start feeding **SWEETLIX® 6% CopperHead® ROL** in early spring before flies begin to appear and continue feeding throughout the year until cold weather restricts fly activity. **SWEETLIX® 6% CopperHead® ROL** may be fed to cattle up until slaughter and to lactating cows.

KEEP OUT OF REACH OF CHILDREN

CAUTION

(See Reverse for First Aid, Precautionary Statements and complete Directions for Use.)

NET WEIGHT: 50 LBS. (22.7 KG)

EPA REG NO. 7698-27

EPA EST. NO. 7698-AL-001

Rabon® is a registered trademark of Bayer HealthCare LLC



7 4 8 9 3 6 0 2 6 9 8 4

Batch #

10897

1118

Manufactured By: **Ridley USA Inc.**
111 West Cherry Street, Suite 500
Mankato, MN 56001

| First Aid | |
|--|---|
| If swallowed | Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin | Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a poison control center or physician for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. For emergencies, call the poison control center at 1-800-222-1222. | |
| NOTE TO PHYSICIANS AND VETERINARIANS: This product is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. Product may cause aspiration pneumonia. Probable mucosal damage may contraindicate the use of gastric lavage. | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

| Personal Protective Equipment ("PPE") |
|--|
| Handlers must wear: long-sleeved shirt and long pants; shoes and socks; and chemical-resistant gloves. |
| User Safety Requirements: Follow manufacturers' instructions for cleaning/maintaining PPE. If no such instructions for washables are provided, use detergent and hot water. Keep and wash PPE separately from other laundry. |
| User Safety Recommendations: Users should thoroughly wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco or using the toilet. Users should immediately remove clothing if pesticide gets inside, then wash thoroughly and put on clean clothing. Users should immediately remove PPE after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. |

Environmental Hazards: This product is highly toxic to fish.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restriction: Not to be used on horses destined for slaughter.

Precautions for Horse Use: This product contains Tetrachlorvinphos, which is a cholinesterase inhibitor. The most frequently reported clinical signs of cholinesterase inhibition in the horse are abdominal pain, lethargy, sweating, tearing and excessive salivation. If these signs are seen in horses, immediately consult your veterinarian. Do not use this product simultaneously or within a week before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. This product has not been tested in breeding horses.

Feeding Instructions

Feed SWEETLIX® 6% CopperHead® ROL continuously on a free choice basis to cattle or horses. Feed as the only free choice source of salt, minerals or vitamins. Feed SWEETLIX® 6% CopperHead® ROL at a level to provide 70 mg RABON® per 100 lbs. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month). Use the table below to determine proper intake levels to deliver the required 70 mg RABON® per 100 lbs. bodyweight per day:

| Animal Weight (lb.) | Supplement Intake (oz.) | Daily Intake of RABON® (lb.) | Daily Intake of RABON® (mg) |
|------------------------|----------------------------|---------------------------------|--------------------------------|
| 500 | 2.5 | 0.15625 | 350 |
| 600 | 3.0 | 0.18750 | 420 |
| 700 | 3.5 | 0.21875 | 490 |
| 800 | 4.0 | 0.25000 | 560 |
| 900 | 4.5 | 0.28125 | 630 |
| 1000 | 5.0 | 0.31250 | 700 |
| 1100 | 5.5 | 0.34375 | 770 |
| 1200 | 6.0 | 0.37500 | 840 |

If animals consume more or less than these levels, feeders should be moved further from or closer to the general loafing or watering areas. If animals still fail to consume the correct amounts, discontinue feeding this free choice supplement. Provide fresh, clean water to all animals at all times. **Follow feeding directions.** Consumption of selenium should not exceed 3 mg per head daily.

Storage and Disposal: Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store in a cool, dry place in the original container. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Nonrefillable container; do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container in a sanitary landfill or by incineration, or if allowed by state or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

For additional information, please contact your SWEETLIX® representative.

NET WEIGHT: 50 LBS. (22.7 KG) Batch #

Manufactured By: **Ridley USA Inc.**
111 West Cherry Street, Suite 500
Mankato, MN 56001

10897
1118

| First Aid | |
|--|---|
| If swallowed | Immediately call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If on skin | Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. |
| If in eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a poison control center or physician for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. For emergencies, call the poison control center at 1-800-222-1222. | |
| NOTE TO PHYSICIANS AND VETERINARIANS: This product is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. Product may cause aspiration pneumonia. Probable mucosal damage may contraindicate the use of gastric lavage. | |

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction to some individuals.

| Personal Protective Equipment ("PPE") |
|--|
| Handlers must wear: long-sleeved shirt and long pants; shoes and socks; and chemical-resistant gloves. |
| User Safety Requirements: Follow manufacturers' instructions for cleaning/maintaining PPE. If no such instructions for washables are provided, use detergent and hot water. Keep and wash PPE separately from other laundry. |
| User Safety Recommendations: Users should thoroughly wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco or using the toilet. Users should immediately remove clothing if pesticide gets inside, then wash thoroughly and put on clean clothing. Users should immediately remove PPE after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. |

Environmental Hazards: This product is highly toxic to fish.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Use Restriction: Not to be used on horses destined for slaughter.

Precautions for Horse Use: This product contains Tetrachlorvinphos, which is a cholinesterase inhibitor. The most frequently reported clinical signs of cholinesterase inhibition in the horse are abdominal pain, lethargy, sweating, tearing and excessive salivation. If these signs are seen in horses, immediately consult your veterinarian. Do not use this product simultaneously or within a week before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. This product has not been tested in breeding horses.

Feeding Instructions

Feed SWEETLIX® 6% CopperHead® ROL continuously on a free choice basis to cattle or horses. Feed as the only free choice source of salt, minerals or vitamins. Feed SWEETLIX® 6% CopperHead® ROL at a level to provide 70 mg RABON® per 100 lbs. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month). Use the table below to determine proper intake levels to deliver the required 70 mg RABON® per 100 lbs. bodyweight per day:

| Animal Weight (lb.) | Supplement Intake (oz.) | Daily Intake of RABON® (lb.) | Daily Intake of RABON® (mg) |
|------------------------|----------------------------|---------------------------------|--------------------------------|
| 500 | 2.5 | 0.15625 | 350 |
| 600 | 3.0 | 0.18750 | 420 |
| 700 | 3.5 | 0.21875 | 490 |
| 800 | 4.0 | 0.25000 | 560 |
| 900 | 4.5 | 0.28125 | 630 |
| 1000 | 5.0 | 0.31250 | 700 |
| 1100 | 5.5 | 0.34375 | 770 |
| 1200 | 6.0 | 0.37500 | 840 |

If animals consume more or less than these levels, feeders should be moved further from or closer to the general loafing or watering areas. If animals still fail to consume the correct amounts, discontinue feeding this free choice supplement. Provide fresh, clean water to all animals at all times. **Follow feeding directions.** Consumption of selenium should not exceed 3 mg per head daily.

Storage and Disposal: Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage:** Store in a cool, dry place in the original container. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Nonrefillable container; do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container in a sanitary landfill or by incineration, or if allowed by state or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

For additional information, please contact your SWEETLIX® representative.

NET WEIGHT: 50 LBS. (22.7 KG) Batch #

Manufactured By: **Ridley USA Inc.**
111 West Cherry Street, Suite 500
Mankato, MN 56001

10897
1118