



SHEEP & GOAT FEED

293040

Pelleted

Formulated to be fed to sheep and goats

MEDICATED

For the prevention of coccidiosis in young sheep caused by *Eimeria bakuensis*, *Eimeria crandalis*, *Eimeria ovinoidalis* and *Eimeria parva*. Feed at least 28 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

ACTIVE DRUG INGREDIENT
Decoquinatate 14 g/ton

Guaranteed Analysis

Crude Protein	(Min)	15.00	%
Crude Fat	(Min)	3.00	%
Crude Fiber	(Max)	12.00	%

Ingredients

Processed Grain By-Products, Grain Products, Roughage Products, Plant Protein Products, Calcium Carbonate, Salt, Potassium Sulfate, Magnesium Sulfate, Kaolin, Vitamin A Supplement, Vitamin D3 Supplement and Vitamin E Supplement.

Feeding Directions

This feed is formulated for sheep and goats and should be fed at a rate of 2% of body weight per head per day.

CAUTION

WARNING: Do not feed to sheep or goats producing milk for food.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed Type C feed or milk replacer at a rate to provide 22.7 mg per 100 lb (0.5 mg/kg) bodyweight. Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Type C feed may be prepared from dry or liquid type B feed.

Always provide a source of fresh clean water, and free choice minerals. When making changes in an animal's diet, allow a 7 to 10 day transition period. Provide ample feeder space for livestock. Do not feed wet, moldy, or insect infested feed

MEDICATED

This feed was designed for mature sheep and goats to help maintain body condition when forages are insufficient. It would be need to be fed during times when nutrient needs are highest such as lactation. It is safe for all classes of sheep and goats other than lactating dairy animals where the milk if to be used for human consumption. This feed is also a good alternative for the performance sheep feed when lambs or kids reach 70 to 80 pounds.