

100 Doses

100 ml

CORYNEBACTERIUM PSEUDOTUBERCULOSIS BACTERIN

For Veterinary Use Only

Indications: For use in healthy goats, three months of age or older, as an aid in the prevention of pyogranulomas/caseous lymphadenitis caused by *Corynebacterium Pseudotuberculosis*.

Directions: SHAKE WELL AND ASEPTICALLY INJECT 1 ML SUBCUTANEOUSLY IN THE SIDE OF THE NECK. Give a second dose 14 days after the first on the opposite side of the neck.

Caution: May cause local swelling. Do not inject into muscle, may cause carcass trim. Do not vaccinate within 60 days of slaughter. Do not contaminate with dirt or chemicals. Use sterile equipment and adequately clean and disinfect bottle cap prior to each entry. Use entire contents when opened. Store at 2° to 7°C. Do not use if emulsion is broken as indicated by separation at the BOTTOM of bottle. In case of anaphylactoid reaction treat symptomatically.

This product license is conditional. Efficacy and potency test studies in progress.

TEXAS VET LAB, INC.
U.S. Veterinary License No. 290
San Angelo, Texas
800-284-8403

**CORYNEBACTERIUM
PSEUDOTUBERCULOSIS
BACTERIN**



Expiration Date:

Serial Number:

PCN: 2A94.00

FOR USE IN GOATS

DOSE: 1 ml SUBCUTANEOUSLY IN SIDE OF NECK

REPEAT IN 14 DAYS ON THE OPPOSITE SIDE OF THE NECK

SEE FRONT PANEL FOR FULL INSTRUCTIONS.